



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## \*KINEMATIC GEOMETRY. INVERSION AND INVERSORS.


By JOHN JAMES QUINN, Ph. D.

This paper is in the nature of a supplement to the one read by the author before Section A, American Association for the Advancement of Science, in Philadelphia, 1904. In that communication two types of Inversors were exhibited and demonstrated. They represented two distinct groups each embodying the property of inversion, and for special cases they become instruments for describing a line which is mathematically straight.

Further investigation has revealed the fact that an infinite variety of those instruments can be made possessing this principle, differing somewhat in appearance from one another yet essentially the same.

The manner in which they can be constructed is set forth in the following theorems:

**THEOREM.** *If from any point P in the axis of symmetry OQ of a concave or convex kite lines be drawn parallel to the shorter sides, and terminated by the longer sides (produced if necessary) the product of the distances  $OP \times OQ$  is constant, whether the point P be taken within or without the points O and Q.*



**GIVEN:** The kite  $ORQV$ ;  $OR$  produced to  $M$ ;  $MP \parallel QV$ ;  $MN$  and  $RS$  perpendicular to  $OQ$ .

**PROOF.**  $OP = ON \pm PN = \sqrt{(OM^2 - MN^2)} \pm \sqrt{(MP^2 - MN^2)}$ ,  
 $OQ = OS \pm QS = \sqrt{(OR^2 - RS^2)} \pm \sqrt{(RQ^2 - RS^2)}$ .

Now  $RS = \frac{RO}{MO} \cdot MN$ ; and  $RQ = \frac{RO}{MO} \cdot MP$ . Substituting we get

$$OP \cdot OQ = \frac{RO}{MO} [OM^2 \pm MP^2], \text{ a constant.}$$


---

\*Read before Section A, American Association for the Advancement of Science, at the New Orleans meeting, December, 1905.

Therefore the points  $O$ ,  $P$ , and  $Q$  are inverse points. Similarly if the points  $P$  and  $Q$  be interchanged. Q. E. D.

SCHOLIUM. From the above it is evident that if the point  $O$  be fixed in position, and the points  $P$ ,  $P'$ , etc., be constrained to move in a circle through  $O$ , the point  $Q$  will describe a straight line, as  $AB$ .

THEOREM. If  $O$  and  $Q$  be two adjacent vertices of a crossed parallelogram  $OQRS$ , and  $P$  be a point situated on a parallel to  $OS$  collinear with  $O$  and  $Q$ , then  $OP \times OQ$  is constant, whether  $P$  be within or without the points  $O$  and  $Q$ .



GIVEN: The crossed parallelogram  $OQRS$ ;  $BA \parallel SO$ ;  $P$  on  $BA$  produced collinear with  $O$  and  $Q$ ;  $AN$  and  $SM$  perpendicular to  $OP$ .

PROOF.  $OP = ON \pm NP = \sqrt{AO^2 - AN^2} \pm \sqrt{AP^2 - AN^2}$ .

$OQ = MQ \pm MO = \sqrt{QS^2 - SM^2} \mp \sqrt{SO^2 - SM^2}$ .

Now  $MS = \frac{OS}{AP} \cdot AN$  and  $QS =$

$\frac{OS}{AP} \cdot AO$ .

Therefore  $OP \cdot OQ = \frac{OS}{AP} [AO^2 \mp AP^2]$  is a constant.

Hence the points  $O$ ,  $P$ , and  $Q$  are inverse points. Q. E. D.

Similarly, is the points  $P'$ ,  $P''$ , etc., be taken.

Evidently then if the point  $O$  be fixed and  $P$  be constrained to move in a circle through  $O$ , the point  $Q$  will move in a straight line, as  $XY$ . The position of the line described by  $Q$  depends upon the position of the center of the circle described by the point  $P$ . It is perpendicular to the line connecting  $O$  to the center of the circle.